



Search List First Prev Next Last

**MicroPatent's Patent Index Database:** [Complete Family of JP7025759A]

11 record(s) found in the family

Order Selected Patent(s)

[no drawing available]

**AU663998B2** **19951026****Title:** (ENG) Chondroprotective agents**Application Number:** AU 6734094 A**Application (Filing) Date:** 19940707**Priority Data:** JP 19418393 19930709 A X;**Inventor(s):** WATANBE KOJU ; NIIMURA KOICHI ; UMEKAWA KIYONORI**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD**IPC (International Class):** A61K03119**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282**Legal Status:** There is no Legal Status information available for this patent

[no drawing available]

**AU6734094A** **19950119****Title:** (ENG) Chondroprotective agents**Application Number:** AU 6734094 D**Application (Filing) Date:** 19940707**Priority Data:** JP 19418393 19930709 A X;**Inventor(s):** WATANBE KOJU ; NIIMURA KOICHI ; UMEKAWA KIYONORI**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD**IPC (International Class):** A61K03119**Legal Status:** There is no Legal Status information available for this patent

[no drawing available]

**CA2126512A1** **19950110****Title:** (ENG) CHONDROPROTECTIVE AGENTS

**Abstract:** A chondroprotective agent comprising a carboxylic acid compound of the general formula (I): (I) wherein R1 and R2 are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

**Application Number:** CA 2126512 A**Application (Filing) Date:** 19940622**Priority Data:** JP 19418393 19930709 A X;**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP**BEST AVAILABLE COPY**

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP

**Last Modification Date:** 20041012

**IPC (International Class):** A61K031215; A61K03119

**Publication Language:** ENG

**Legal Status:**

Date +/- Code Description

20030403 (+) EEER EXAMINATION REQUEST Effective date: 19940622;

20030403 (-) FZDEDEAD Effective date: 19990622;

[no drawing available]

**CN1100631A**  **19950329**

**Title:** (ENG) Chondroprotective agents

**Application Number:** CN 94108231 A

**Application (Filing) Date:** 19940708

**Priority Data:** JP 19418393 19930709 A X;

**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP

**Last Modification Date:** 20050208

**IPC (International Class):** A61K03119

**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282

**Legal Status:** There is no Legal Status information available for this patent

**DE69410024D1**  **19980610** [FullText](#)

**Title:** (GER) Knorpelschuetzende Arzneimittel

**Application Number:** DE 69410024 A

**Application (Filing) Date:** 19940627

**Priority Data:** JP 19418393 19930709 A X;

**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP

**IPC (International Class):** A61K03119; A61K031215

**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282

**Legal Status:**

Date +/- Code Description

19990602 (+) 8364 NO OPPOSITION DURING TERM OF OPPOSITION

20010809 (-) 8339 CEASED/NON-PAYMENT OF THE ANNUAL FEE

**DE69410024T2**  **19981119** [FullText](#)

**Title:** (GER) Knorpelschuetzende Arzneimittel

**Application Number:** DE 69410024 T

**Application (Filing) Date:** 19940627

**Priority Data:** JP 19418393 19930709 A X;

**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP

**IPC (International Class):** A61K03119; A61K031215

**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282

**Legal Status:**

Date +/- Code Description

19990602 (+) 8364 NO OPPOSITION DURING TERM OF OPPOSITION


20010809 (-) 8339 CEASED/NON-PAYMENT OF THE ANNUAL FEE

EPO Register

**EP0633021A1**  **19950111** [FullText](#)

**Title:** (ENG) Chondroprotective agents**Abstract:** (ENG)

A chondroprotective agent comprising a carboxylic acid compound of the general formula (I):



wherein R<sup>1</sup> and R<sup>2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

**Application Number:** EP 94109873 A**Application (Filing) Date:** 19940627**Priority Data:** JP 19418393 19930709 A I;**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP**IPC (International Class):** A61K03119; A61K031215**ECLA (European Class):** A61K03119; A61K031215**Designated Countries:**

- Designated States: CH DE FR GB IT LI SE

**Publication Language:** ENG**Agent(s):** Minderop, Ralph H., Dr. rer. nat. et al 00073451 Cohausz & Florack Patentanwalte Postfach 33 02 29 40435 Duesseldorf DE**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282**Other Abstracts for This Document:** CAN122(09)096508H; DERC95-038282**Non-Patent Citations:**

- INDIAN J. EXP. BIOL., vol.25, no.3, March 1987 pages 187 - 189 A.S.CHAWLA ET AL. 'Anti-Inflammatory Action of Ferulic Acid and Its Esters in Carrageenan Induced Rat Paw Oedema Model'
- ACTA ZOOL. SIN., vol.38, no.1, March 1992 pages 106 - 107 B.- S. ZHOU ET AL. 'ULTRASTRUCTURAL OBSERVATION ON THE EFFECTS OF FERULIC ACID ON THE CARTILAGE PROTEOGLYCANS IN YOUNG RATS'
- WUHAN DAXUE XUEBAO, ZIRAN KEXUEBAN, no.4, 1990 pages 93 - 97 B.ZHOU ET AL. 'ACTION OF FERULIC ACID AND SODIUM SELEMITATE ON YOUNG RATS'
- BIOTECHNOL. THER., vol.3, no.3-4, 1992 pages 119 - 135 B.A. 'T HART ET AL. 'METABOLIC ACTIVATION OF PHENOLS BY STIMULATED NEUTROPHILS: A CONCEPT FOR A SELECTIVE TYPE OF ANTI-INFLAMMATORY DRUG'
- ACUPUNCT. ELECTRO-THER. RES. INT. J., vol.7, no.2-3, 1982 pages 173 - 202 J. CYONG ET AL. 'A PHARMACOLOGICAL STUDY OF THE ANTI-INFLAMMATORY ACTIVITY OF CHINESE HERBS. A REVIEW'
- CLIN. CHIM. ACTA, vol.73, no.1, 1976 pages 121 - 125 M.R. WILLS ET AL. 'THE EFFECT OF URAEMIC METABOLITES ON PARATHYROID EXTRACT-INDUCED BONE RESORPTION IN VITRO'

**Legal Status:**

Date	+/- Code	Description
19950111(+)	17P	REQUEST FOR EXAMINATION FILED Effective date: 19940627;
19950111(+)	AK	DESIGNATED CONTRACTING STATES: Kind code of corresponding patent document: A1; CH DE FR GB IT LI SE
19960918(+)	17Q	FIRST EXAMINATION REPORT Effective date: 19960805;
19980506(+)	AK	DESIGNATED CONTRACTING STATES: Kind code of corresponding patent document: B1; CH DE FR GB IT LI SE
19980515	REG	REFERENCE TO A NATIONAL CODE : CH; : EP;
19980610	REF	CORRESPONDS TO: Corresponding patent document:

69410024; Country code of corresponding patent document: DE; Publication date of corresponding patent document: 19980610;

19980703(+)ITF IT: TRANSLATION FOR A EP PATENT FILED New owner name: MODIANO & ASSOCIATI S.R.L.;

19980710(+)ET FR: TRANSLATION FILED

19981028 RAP2 PATENT OWNER REASSIGNMENT (CORRECTION) New owner name: KUREHA CHEMICAL INDUSTRY CO., LTD.;

19981104 RAP2 PATENT OWNER REASSIGNMENT (CORRECTION) New owner name: KUREHA CHEMICAL INDUSTRY CO., LTD.;

19981113 REG REFERENCE TO A NATIONAL CODE : CH; : PL;

19990428(+)26N NO OPPOSITION FILED

20001213 (-) 25 LAPSED IN A CONTRACTING STATE : CH; Effective date: 19980811;

20001213 (-) 25 LAPSED IN A CONTRACTING STATE : LI; Effective date: 19980811;

20001213 (-) 25 LAPSED IN A CONTRACTING STATE : SE; Effective date: 19980806;

20001227 (-) R25 LAPSED AS TO RULE 92 1 P (CORRECTION) : CH; Effective date: 19980506;

20001227 (-) R25 LAPSED AS TO RULE 92 1 P (CORRECTION) : LI; Effective date: 19980506;


20001227 (-) R25 LAPSED AS TO RULE 92 1 P (CORRECTION) : SE; Effective date: 19980806;

20010214 (-) GBPCGB: EUROPEAN PATENT CEASED THROUGH NON-PAYMENT OF RENEWAL FEE Effective date: 20000627;

20010330 REG REFERENCE TO A NATIONAL CODE : FR; : ST;

**Additional Information:**

- Date of request for examination 19940627
- Date of dispatch of first examination report 19960805
- Patent bulletin/ gazette information 20001227 200052
- Lapse of a Patent , 19980506 CH, 19980506 LI, 19980806 SE

**EPO Register****EP0633021B1**  **19980506** [FullText](#)**Title:** (ENG) Chondroprotective agents**Abstract:** (ENG)

A chondroprotective agent comprising a carboxylic acid compound of the general formula (I):

wherein R<sup>1</sup> and R<sup>2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

**Application Number:** EP 94109873 A**Application (Filing) Date:** 19940627**Priority Data:** JP 19418393 19930709 A I;**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP**IPC (International Class):** A61K03119; A61K031215**ECLA (European Class):** A61K03119; A61K031215**Designated Countries:**

- Designated States: CH DE FR GB IT LI SE

**Publication Language:** ENG**Agent(s):** Minderop, Ralph H., Dr. rer. nat. et al 00073451 Cohausz & Florack Patentanwalte Postfach 33 02 29 40435 Duesseldorf DE

**Other Abstracts for Family Members:** CHEMABS122(09)096508H;  
DERABS C95-038282

**Non-Patent Citations:**

- INDIAN J. EXP. BIOL., vol.25, no.3, March 1987 pages 187 - 189 A.S.CHAWLA ET AL. 'Anti-Inflammatory Action of Ferulic Acid and Its Esters in Carrageenan Induced Rat Paw Oedema Model'
- ACTA ZOOL. SIN., vol.38, no.1, March 1992 pages 106 - 107 B.- S. ZHOU ET AL. 'ULTRASTRUCTURAL OBSERVATION ON THE EFFECTS OF FERULIC ACID ON THE CARTILAGE PROTEOGLYCANS IN YOUNG RATS'
- WUHAN DAXUE XUEBAO, ZIRAN KEXUEBAN, no.4, 1990 pages 93 - 97 B.ZHOU ET AL. 'ACTION OF FERULIC ACID AND SODIUM SELEMITE ON YOUNG RATS'
- BIOTECHNOL.THER., vol.3, no.3-4, 1992 pages 119 - 135 B.A. 'T HART ET AL. 'METABOLIC ACTIVATION OF PHENOLS BY STIMULATED NEUTROPHILS: A CONCEPT FOR A SELECTIVE TYPE OF ANTI-INFLAMMATORY DRUG'
- ACUPUNCT. ELECTRO-THER. RES. INT. J., vol.7, no.2-3, 1982 pages 173 - 202 J.CYONG ET AL. 'A PHARMACOLOGICAL STUDY OF THE ANTI-INFLAMMATORY ACTIVITY OF CHINESE HERBS. A REVIEW'
- CLIN.CHIM.ACTA, vol.73, no.1, 1976 pages 121 - 125 M.R.WILLS ET AL. 'THE EFFECT OF URAEMIC METABOLITES ON PARATHYROID EXTRACT-INDUCED BONE RESORPTION IN VITRO'

**Legal Status:**

Date	+/-	Code	Description
19950111	(+)	17P	REQUEST FOR EXAMINATION FILED Effective date: 19940627;
19950111	(+)	AK	DESIGNATED CONTRACTING STATES: Kind code of corresponding patent document: A1; CH DE FR GB IT LI SE
19960918	(+)	17Q	FIRST EXAMINATION REPORT Effective date: 19960805;
19980506	(+)	AK	DESIGNATED CONTRACTING STATES: Kind code of corresponding patent document: B1; CH DE FR GB IT LI SE
19980515		REG	REFERENCE TO A NATIONAL CODE : CH; : EP;
19980610		REF	CORRESPONDS TO: Corresponding patent document: 69410024; Country code of corresponding patent document: DE; Publication date of corresponding patent document: 19980610;
19980703	(+)	ITF	IT: TRANSLATION FOR A EP PATENT FILED New owner name: MODIANO & ASSOCIATI S.R.L.;
19980710	(+)	ET	FR: TRANSLATION FILED
19981028		RAP2	PATENT OWNER REASSIGNMENT (CORRECTION) New owner name: KUREHA CHEMICAL INDUSTRY CO., LTD.;
19981104		RAP2	PATENT OWNER REASSIGNMENT (CORRECTION) New owner name: KUREHA CHEMICAL INDUSTRY CO., LTD.;
19981113		REG	REFERENCE TO A NATIONAL CODE : CH; : PL;
19990428	(+)	26N	NO OPPOSITION FILED
20001213	(-)	25	LAPSED IN A CONTRACTING STATE : CH; Effective date: 19980811;
20001213	(-)	25	LAPSED IN A CONTRACTING STATE : LI; Effective date: 19980811;
20001213	(-)	25	LAPSED IN A CONTRACTING STATE : SE; Effective date: 19980806;
20001227	(-)	R25	LAPSED AS TO RULE 92 1 P (CORRECTION) : CH; Effective date: 19980506;
20001227	(-)	R25	LAPSED AS TO RULE 92 1 P (CORRECTION) : LI; Effective date: 19980506;
20001227	(-)	R25	LAPSED AS TO RULE 92 1 P (CORRECTION) : SE; Effective date: 19980806;
20010214	(-)	GBPCGB	EUROPEAN PATENT CEASED THROUGH NON-PAYMENT OF RENEWAL FEE Effective date: 20000627;
20010330		REG	REFERENCE TO A NATIONAL CODE : FR; : ST;

**Additional Information:**

- Date of request for examination 19940627
- Date of dispatch of first examination report 19960805

- Patent bulletin/ gazette information 20001227 200052
- Lapse of a Patent , 19980506 CH, 19980506 LI, 19980806 SE

[no drawing available]

**JP3490114B2**  **20040126** [FullText](#)

**Application Number:** JP 19418393 A

**Application (Filing) Date:** 19930709

**Priority Data:** JP 19418393 19930709 A X;

**IPC (International Class):** A61K03119; A61K031215; A61P01902; A61P02900

**Legal Status:** There is no Legal Status information available for this patent


**JP7025759A**  **19950127** [FullText](#)

**Title:** (ENG) AGENT FOR PROTECTING CARTILAGE

**Abstract:** (ENG)

**PURPOSE:** To obtain the agent composed of a carboxylic acid having a specific structure, exhibiting cartilage- protecting action by its strong action to suppress the decrease of glycosaminoglycan constituting cartilage matrix, having low toxicity and useful for the treatment of rheumatoid arthritis, etc.

**CONSTITUTION:** This agent is composed of a carboxylic acid of the formula (R1 and R2 are H or 1-6C alkyl), its cis or trans isomer, its salt or its ester. The compound of the formula is e.g. 3-ethoxy-4-hydroxycinnamic acid obtained by reacting 3-ethoxy-4-hydroxybenzaldehyde with malonic acid in the presence of a base such as pyridine. The amount of the active component in the agent is preferably 0.1-70wt.%. The agent is administered preferably at a daily rate of 0.5-200mg/kg (body weight) usually in one dose or 2-4 divided doses.



**Application Number:** JP 19418393 A

**Application (Filing) Date:** 19930709

**Priority Data:** JP 19418393 19930709 A X;

**Inventor(s):** NIIMURA KOICHI ; UMEKAWA KIYONORI ; WATANABE TAKATOSHI

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD

**IPC (International Class):** A61K03119; A61K031215

**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C95-038282

**Legal Status:** There is no Legal Status information available for this patent

**US5621009A**  **19970415** [FullText](#)

**Title:** (ENG) Chondroprotective agents

**Abstract:** (ENG)

A chondroprotective agent comprising a carboxylic acid compound of the general formula (I): STR1 wherein R<sup>sup.1</sup> and R<sup>sup.2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

**Application Number:** US 53551695 A

**Application (Filing) Date:** 19950928

**Priority Data:** US 53551695 19950928 A I; JP 19418393 19930709 A R; US 27257494 19940711 A I;

**Related Application(s):** 08/272574 19940711 ABANDONED

**Inventor(s):** WATANABE KOJU JP ; NIIMURA KOICHI JP ; UMEKAWA KIYONORI JP

**Assignee/Applicant/Grantee:** KUREHA CHEMICAL IND CO LTD JP

**Last Modification Date:** 20050627

**IPC (International Class):** A61K03119

**ECLA (European Class):** A61K0031216**US Class:** 514568; 514825**Agent(s):** Sughrue, Mion, Zinn, Macpeak & Seas**Examiner Primary:** Criares, Theodore J.**Other Abstracts for Family Members:** CHEMABS122(09)096508H; DERABS C1995-038282**Non-Patent Citations:**

- Indian J. Exp. Biol., "Anti-Inflammatory Action of Ferulic Acid and Its Esters in Carrageenan Induced Rat Paw Oedema Model" by A Chawla et al, vol. 25, No. 3, Mar. 1987, pp. 187-189.
- Acta Zool. Sin., "Ultrastructural Observation on the Effects of Ferulic Acid on the Cartilage Proteoglycans in young Rats" by B Zhou et al, vol. 38, No. 1, Mar. 1992, pp. 106-107.
- Wuhan Daxue Xuebao, Ziran Kexueban, "Action of Ferulic Acid and Sodium Selenite on Young Rats" by B Zhou et al, No. 4, 1990, pp. 93-97.
- Biotechnol. Ther., "Metabolic Activation of Phenols by Stimulated Neutrophils: A Concept for a Selective Type of Anti-Inflammatory Drug" by B T Hart et al, vol. 3, No. 3-4, 1992, pp. 119-135.
- Acupunct. Electro-Ther. Res. Int. J., "A Pharmacological Study of the Anti-Inflammatory Activity of Chinese Herbs. A Review." by J. Cyong et al, vol. 7, No. 2-3, 1982, pp. 173-202.
- Clin. Chim. Act., "The Effect of Uraemic Metabolites on Parathyroid Extract-Induced Bone Resorption In Vitro" by M Wills et al, vol. 73, No. 1, 1976, pp. 121-125.

**Patents Cited:**

→ EP0083285 19830700

**Legal Status:**

Date +/- Code Description

20050614 (-) FP EXPIRED DUE TO FAILURE TO PAY MAINTENANCE FEE Effective date: 20050415;

**US Post Issuance:**

- US Expiration Date: 20050415 20050614 DUE TO FAILURE TO PAY MAINTENANCE FEES

**Additional Information:**

- Number of claims 3
- Exemplary claim number(s) 1
- National classifications searched , 514/568



Search



List



First



Prev



Next



Last

Copyright © 2002, MicroPatent, LLC. The contents of this page are the property of MicroPatent LLC including without limitation all text, html, asp, javascript and xml. All rights herein are reserved to the owner and this page cannot be reproduced without the express permission of the owner.



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number: **0 633 021 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 94109873.3

(51) Int. Cl.<sup>6</sup>: A61K 31/19, A61K 31/215

(22) Date of filing: 27.06.94

(30) Priority: 09.07.93 JP 194183/93

(43) Date of publication of application:  
11.01.95 Bulletin 95/02

(84) Designated Contracting States:  
CH DE FR GB IT LI SE

(71) Applicant: KUREHA CHEMICAL INDUSTRY CO.,  
LTD.  
9-11, Horidome-cho, 1-chome  
Nihonbashi  
Chuo-ku  
Tokyo 103 (JP)

(72) Inventor: Watanabe, Koju  
2-5-7, Tsurumai  
Sakado-shi,

Saitama 350-02 (JP)

Inventor: Nilmura, Koichi

Rune Warabi 1-718,

1-17-30, Chuo

Warabi-shi,

Saitama 335 (JP)

Inventor: Umekawa, Kiyonori

5-4-309, Mihama

Urayasu-shi,

Chiba 279 (JP)

(74) Representative: Minderop, Ralph H. Dr. rer.nat.  
et al

Cohausz & Florack


Patentanwälte

Bergiusstrasse 2 b

D-30655 Hannover (DE)

(54) Chondroprotective agents.

(57) A chondroprotective agent comprising a carboxylic acid compound of the general formula (I):



wherein R<sup>1</sup> and R<sup>2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

EP 0 633 021 A1

## BACKGROUND OF THE INVENTION

## 1. Field of the Invention

5 The present invention relates to an agent for protecting cartilage, i.e., a chondroprotective agent, more particularly, a chondroprotective agent containing a carboxylic acid compound of the general formula (I) as mentioned below or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof.

## 2. Description of the Related Art

10 There are various types of arthropathy, for example, rheumatoid arthritis, rheumatic fever, and osteoarthritis. Many people particularly suffer from rheumatoid arthritis and osteoarthritis. These diseases have been studied as the major types of arthropathy. There are congenital and secondary osteoarthritis, and further primary osteoarthritis caused by degeneration of the articular cartilage along with aging. Patients  
15 suffering from primary osteoarthritis have recently been increasing along with the increase in the population of the aged.

Although there are considerable differences of the causes and conditions between rheumatoid arthritis and osteoarthritis, the articular function becomes eventually obstructed by the destruction of the cartilage in both of rheumatoid arthritis and osteoarthritis.

20 The first choice of medicines for the treatment of rheumatic diseases such as rheumatoid arthritis, rheumatic fever, systemic lupus erythematosus, and osteoarthritis are analgesic and anti-inflammatory agents, for example, aspirin or indometacin. Further, gold compounds such as Shiosol, immunomodulators, steroids, or D-penicillamine are used as medicines for the treatment of rheumatoid arthritis.

The above conventional analgesic and anti-inflammatory agents, however, were not effective against the  
25 destruction of the articular cartilage, and in fact, sometimes exhibited adverse effect in experiments using chondrocytes. Further, no inhibitory effect on the destruction of articular cartilage was observed in the above medicines for the treatment of rheumatoid arthritis.

It is known that caffeic, ferulic, isoferulic, or 3-ethoxy-4-hydroxycinnamic acid may be used in the following pharmaceutical applications: Caffeic and ferulic acids exhibit an antiviral activity [Japanese  
30 Unexamined Patent Publication (Kokai) No. 4-234319]. Caffeic acid may be used as a calcium antagonist [Japanese Unexamined Patent Publication (Kokai) No. 4-243822], an anti-allergenic agent [Japanese Unexamined Patent Publication (Kokai) No. 59-155314], and the like. Caffeic acid and methyl caffeate can be used for the treatment of autoimmune disorders (WO91/17749). Further, ferulic and isoferulic acids may be used as an antitumor agent for preventing drug resistance in immunochemotherapy [Japanese Unex-  
35 amined Patent Publication (Kokai) No. 56-115716]. Further, ferulic acid may be used as a composition for mitigating hyperlipemia and platelet aggregation (U.S. Patent No. 4,842,859). Further, ferulic acid and esters thereof exhibit an anti-inflammatory action (A.S. Chawla et al., Indian J. Exp. Biol., Vol. 25, No. 3, pp. 187 to 189, 1987). 3-Ethoxy-4-hydroxycinnamic acid may be used as an ultraviolet absorbent for cosmetics [Japanese Unexamined Patent Publication (Kokai) No. 64-13018].

40 The above carboxylic acids, the cis- and trans-isomers thereof, the salts and the esters have not, however, been known to be useful as chondroprotective agents.


## SUMMARY OF THE INVENTION

45 The present inventors engaged in intensive research to develop a chondroprotective agent for suppressing the destruction of the articular cartilage and as a result found that the particular carboxylic acid compounds, and cis- and trans-isomers thereof, and pharmaceutically acceptable salts, and esters thereof showed significant inhibition of the depletion of proteoglycan which is a major component of the cartilage matrix, and therefore, are useful as a chondroprotective agent for prohibiting the destruction of the articular  
50 cartilage.

Accordingly, the object of the present invention is to provide a chondroprotective agent containing as an active ingredient a particular carboxylic acid compound of the general formula (I), or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof.

Other objects and effects of the present invention will be clear from the following description.

55 The present invention relates to a chondroprotective agent comprising a carboxylic acid compound of the general formula (I):



wherein R<sup>1</sup> and R<sup>2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof (hereinafter referred to as "the present substance").

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

in the general formula (I), R<sup>1</sup> and R<sup>2</sup> are preferably a hydrogen atom or lower alkyl groups having 1 to 4 carbon atoms, more preferably methyl, ethyl, n-propyl, or n-butyl groups.

There exist a cis-form and a trans-form as to the configuration of the phenyl group and carbonyl group bonded at both sides of the double bond in the present substance. Each of the cis- and trans-forms may be used for the chondroprotective agent of the present invention.

As the pharmaceutically acceptable salts, there may be mentioned salts with alkali metals (such as sodium or potassium), salts with alkali earth metals (such as calcium or magnesium), aluminum salts, ammonium salts, salts with various amine compounds (such as primary, secondary, or tertiary amines), or the like.

The esters may be those with aromatic or aliphatic alcohols, preferably those with aliphatic alcohols, more preferably those with aliphatic alcohols having 1 to 6 carbon atoms. As even more preferable examples, there may be mentioned methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, or isopentyl esters.

Examples of the present substance are listed in Table 1. Table 1 shows only the trans-forms, to which, however, the present invention is not limited.

Table 1

R <sup>1</sup>	R <sup>2</sup>	Ester	Name	Abbreviation
H	H		Caffeic acid	CFA
Me	H		Ferulic acid	FLA
H	Me		Isoferulic acid	IFLA
Et	H		3-Ethoxy-4-hydroxycinnamic acid	3-EtO-4-HO-CA
Me	H	(Na salt)	Sodium ferulate	NaFLA
Me	H	Me	Methyl ferulate	MeFLA
Me	H	Et	Ethyl ferulate	EtFLA
Me	H	nPr	n-Propyl ferulate	nPrFLA
Me	H	nBu	n-Butyl ferulate	nBuFLA
Me: methyl, Et: ethyl, nPr: n-propyl, nBu: n-butyl				

Many carboxylic acid compounds described in Table 1 are commercially available. For example, caffeic acid is available from Tokyo Kasei Kogyo Co. and ferulic and isoferulic acids are available from Sigma Chemical Co. Further, 3-ethoxy-4-hydroxycinnamic acid may be prepared by reacting 3-ethoxy-4-hydroxybenzaldehyde and malonic acid in the presence of a base such as pyridine (see Example 1 as below).

The pharmaceutically acceptable salts of the free carboxylic acids of the general formula (I) may be prepared by reacting hydroxides or carbonates of alkali metals, alkaline earth metals or ammonium, with an equimolar amount of said free carboxylic acids in a solvent such as water.

The ester of the free carboxylic acid of the general formula (I) may be prepared by reacting alcohol compounds with the above-mentioned free carboxylic acids in one of the following methods comprising:

(1) reacting an alcohol compound and the carboxylic acid in a suitable solvent in the presence of an acid catalyst, for example, an inorganic acid, such as hydrochloric, sulfuric or phosphoric acid, or an organic acid, such as acetic or p-toluene sulfonic acid;

(2) reacting an alcohol compound and the carboxylic acid in the presence of a condensation agent, such as dicyclohexylcarbodiimide, N,N'-carbonyldi(2-methyl-imidazole), diphenylketene-N-cyclohexylimine, alkoxyacetylene, ethyl polyphosphate, thionyl chloride, or oxalyl chloride, in an organic solvent, such as dimethylformamide, acetone, dioxane, acetonitrile, chloroform, ethylene chloride, tetrahydrofuran, or pyridine, generally with cooling or at room temperature;

(3) reacting a carboxylic acid anhydride and an alcohol compound in the presence of a basic material, such as triethylamine, pyridine, or methylethylpyridine; or

(4) reacting a carboxylic acid halide, for example, acyl halide, such as chlorides or bromides, and an alcohol compound in a solvent containing a basic substance, such as triethylamine, pyridine or methylethylpyridine, or in a basic solvent, such as pyridine.

As clear from the toxicity shown below, the present substance is an extremely safe compound.

#### (1) Acute toxicity of caffeic acid

The LDLO (minimum lethal dose) for intraperitoneal administration in rats is 1500 mg/kg (Toxicology and Applied Pharmacology, Vol. 36, p. 227, 1976).

#### (2) Acute toxicity of ferulic, isoferulic, and 3-ethoxy-4-hydroxycinnamic acids

Each of the above acids was orally administered at the dose of 400 mg/kg to ddY male mice (five mice per group) and the mice were observed for 1 week after the administration. The substances to be tested were orally administered after dispersed in an aqueous solution of 0.3 % CMC (carboxymethylcellulose) and 0.05 % Tween 80 (polyoxyethylene sorbitan monooleate; Tokyo Kasei Kogyo Co.). No deaths or no abnormalities in the body weight or general condition were observed in each of the substances to be tested.

The same results were obtained where sodium ferulate, methyl ferulate, ethyl ferulate, propyl ferulate, or butyl ferulate was administered.

The LD<sub>50</sub> values of methyl ferulate and isopropyl ferulate in intraperitoneal administration to mice are 1188.5 mg/kg and 1059.2 mg/kg, respectively (A.S. Chawla et al., Indian J. Exp. Biol., Vol. 25, No. 3, pp. 187 to 189, 1987).

As a pharmacological effect, the present substance exhibits the function to inhibit destruction of chondrocyte matrix in cultured chondrocytes (derived from cartilage of rabbit shoulder and knee joints) (see Example 2 as below). Although the ester compound of the present invention does not always exhibit such a pharmacological effect in vitro, the function to inhibit cartilage destruction induced by granuloma was observed in animal experiments of a mouse air pouch model.

Accordingly, the present substance is useful as a chondroprotective agent for treating various types of arthropathy accompanying the cartilage destruction of joints. Examples of such arthropathy are rheumatoid arthritis, osteoarthritis, peri arthritis humeroscapularis, shoulder-arm-neck syndrome, lumbago, etc.

The chondroprotective agent containing the present substance as an active ingredient may be in the form of any conventional formulation. The chondroprotective agent may contain the present substance alone, or a mixture of the present substance with any pharmaceutically acceptable carrier or diluent. The chondroprotective agent may contain the active ingredient in an amount of 0.01 to 100 percent by weight, preferably 0.1 to 70 percent by weight.

The chondroprotective agent of the present invention may be administered orally or by some other routes.

The dose of the chondroprotective agent according to the present invention varies with the patient (animal or human), age, individual differences, state of illness, and the like. Generally speaking, however, when a human is treated, the dose of oral administration of the present substance is in the range of 0.1 to 500 mg/kg (body weight) per day, preferably 0.5 to 200 mg/kg (body weight), which is usually divided into 1 to 4 dosages in a day, although the dose outside the above range may sometimes be administered.

#### EXAMPLE

The present invention now will be further illustrated by, but is by no means limited to, the following Examples.

Example 1: Preparation of Present Substance

## (1) Preparation of 3-ethoxy-4-hydroxycinnamic acid

- 5 Malonic acid (1.56 g) and pyridine (3.0 ml) were placed in an egg-plant type flask (100 ml). Further, 3-ethoxy-4-hydroxybenzaldehyde (1.66 g) and aniline (30  $\mu$ l) were added thereto and a reaction was performed at 55 °C for 3 hours. The reaction solution was acidified by adding dilute sulfuric acid. The crude crystals were obtained by filtration, washed with water, and then dried to obtain the above-identified compound (1.1 g, 56.1%) as a light yellowish crystal.
- 10 Melting point: 157-158 °C  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>,  $\delta$ ppm): 1.40 (t, 3H, J = 6.87 Hz, CH<sub>3</sub>), 4.19 (q, 2H, J = 6.87 Hz, CH<sub>2</sub>), 6.36 (d, 1H, J = 16.04 Hz), 6.87 (d, 1H, J = 8.2 Hz), 7.13 (dd, 1H, J = 8.2, 1.8 Hz), 7.31 (s, 1H), 7.59 (d, 1H, J = 16.04 Hz)
- IR spectrum (cm<sup>-1</sup>): 3550s, 2980s, 2900s, 2830s, 2700m, 2600m, 2550m, 2510m, 1675s, 1625s, 1590s,  
 1515s, 1482m, 1435m, 1415m, 1330m, 1290s, 1260s, 1235s, 1195s, 1165s, 1121s, 1035s

## (2) Preparation of methyl ferulate

- 20 Ferulic acid (4.85 g) and then methyl alcohol (50 ml) containing hydrogen chloride (4%) were placed in an egg-plant type flask (200 ml), and heated under reflux for 1 hour. The disappearance of the starting material was confirmed by thin layer chromatography (TLC: n-hexane/ethyl acetate = 2/1), then the solvent was evaporated under reduced pressure by a rotary evaporator. Benzene was added to the residue, and then, the solvent was again evaporated by a rotary evaporator to obtain the crude product. The resulting crude product was separated and purified by silica gel column chromatography (diameter = 5.5 x 7.0 cm;  
 25 60 g; n-hexane/ethyl acetate = 2/1) to obtain the above-identified compound. The product was crystallized from n-hexane to obtain a white crystal (5.09 g, 97.9%).
- Melting point: 63-64 °C  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>,  $\delta$  ppm): 3.79 (s, 3H, COOCH<sub>3</sub>), 3.90 (s, 3H, OCH<sub>3</sub>), 6.08 (s, 1H, OH), 6.28 (d, 1H, J = 15.6 Hz, H3), 6.91 (d, 1H, J = 8.25 Hz), 7.01 (d, 1H, J = 1.74 Hz), 7.04 (dd, 1H, J = 8.25, 1.74 Hz), 7.62 (d, 1H, J = 15.6 Hz, H4)

## (3) Preparation of ethyl ferulate

- 35 Ferulic acid (4.85 g) and then ethyl alcohol (50 ml) containing hydrogen chloride (4%) were placed in an egg-plant type flask (200 ml), and heated under reflux for 2.5 hours. The disappearance of the starting material was confirmed by TLC (n-hexane/ethyl acetate = 2/1), then the solvent was evaporated under reduced pressure by a rotary evaporator. Benzene was added to the residue, and then, the solvent was again evaporated by a rotary evaporator to obtain the wine-colored crude product (5.79 g). The resulting crude product was separated and purified by silica gel column chromatography (diameter = 5.5 x 7.0 cm;  
 40 60 g; n-hexane/ethyl acetate = 3/1) to obtain the above-identified compound (5.06 g; 91.2%). The product was crystallized from n-hexane to obtain a white crystal (4.743 g, 85.3%).
- Melting point: 41-43 °C  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>,  $\delta$  ppm): 1.33 (t, 3H, COOCH<sub>2</sub>CH<sub>3</sub>), 3.91 (s, 3H OCH<sub>3</sub>), 4.25 (q, 2H, J = 7.33 Hz), 6.06 (s, 1H, OH), 6.28 (d, 1H, J = 15.6 Hz), 6.91 (d, 1H, J = 6.8 Hz), 7.02 (s, 1H), 7.06 (d, 1H, J = 8.25 Hz), 7.61 (d, 1H, J = 15.6 Hz)

## (4) Preparation of n-propyl ferulate

- 50 Ferulic acid (4.85 g) and then n-propyl alcohol (50 ml) containing hydrogen chloride (4%) were placed in an egg-plant type flask (200 ml), and heated under reflux overnight. The disappearance of the starting material was confirmed by thin layer chromatography (TLC: n-hexane/ethyl acetate = 2/1), then the solvent was evaporated under reduced pressure by a rotary evaporator. Benzene was added to the residue, and then, the solvent was again evaporated by a rotary evaporator to obtain the crude product. The resulting crude product was separated and purified by silica gel column chromatography (diameter = 5.5 x 7.0 cm;  
 55 60 g; n-hexane/ethyl acetate = 2/1) to obtain the above-identified compound (5.48 g, 91.6%) as colorless oil.
- <sup>1</sup>H-NMR (CDCl<sub>3</sub>,  $\delta$  ppm): 0.99 (t, 3H, J = 7.33 Hz), 1.72 (s, 2H, J = 7.33 Hz), 3.90 (s, 3H, OCH<sub>3</sub>), 4.16 (t, 2H, J = 6.9 Hz), 6.11 (s, 1H, OH), 6.30 (d, 1H, J = 15.6 Hz), 6.91 (d, 1H, J = 8.25 Hz), 7.02 (s, 1H), 7.05

(dd, J = 8.23 Hz), 7.61 (d, 1H, J = 15.6 Hz)

#### (5) Preparation of n-butyl ferulate

5 Ferulic acid (5 g) and then n-butyl alcohol (50 ml) containing hydrogen chloride (4%) were placed in an egg-plant type flask (200 ml), and heated under reflux for about 2 hours. The disappearance of the starting material was confirmed by thin layer chromatography (TLC: n-hexane/ethyl acetate = 2/1), then the solvent was evaporated under reduced pressure by a rotary evaporator. Benzene was added to the residue, and then, the solvent was again evaporated by a rotary evaporator to obtain the crude product. The resulting  
10 crude product was separated and purified by silica gel column chromatography (diameter = 5.5 x 8.5 cm; 70 g; n-hexane/ethyl acetate = 2/1) to obtain the above-identified compound (6.25 g, 99%) as colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ ppm): 0.96 (t, 3H, J = 7.33 Hz), 1.44 (seq, 2H J = 7.33 Hz), 1.68 (quint, 2H, J = 6.88 Hz), 3.91 (s, 3H), 4.21 (t, 2H, J = 6.88 Hz), 6.04 (s, 1H, OH), 6.29 (d, J = 16.0 Hz, 1H), 6.91 (d, J = 8.25 Hz), 7.02 (s, 1H), 7.05 (dd, J = 8.25 Hz)

15

#### Example 2: Effect of Test Compounds on Proteoglycan Depletion in Chondrocyte Culture

##### (a) Preparation of Cultured Chondrocytes

20 The cartilages were sterily extracted from the shoulder and knee joints of rabbits (New Zealand White Rabbits) (body weight of 1 to 1.5 kg). The cartilages were thoroughly washed with phosphate buffer (PBS (-): free of Ca<sup>2+</sup>, Mg<sup>2+</sup>), Hanks' solution and 0.1% EDTA-PBS (-), and then cut into small segments (1 mm x 1 mm x 1mm). After PBS (-) containing 0.1% EDTA was added, the segments were allowed to stand in an incubator of 37°C for 30 minutes. Then, the segments were treated with a trypsin solution (0.25%) at 37°C  
25 for one hour to remove the connective tissue adhered to the cartilage. After the supernatant had been removed, the cartilages were treated for about 2 to 2.5 hours in a Ham F-12 medium containing 10 % fetal bovine serum (FBS) and 0.2 % collagenase. Then, the collagenase solution was centrifuged (1500 r.p.m.), and the residual chondrocytes were washed twice with a Ham F-12 medium (chondrocyte culture medium) containing 10 % FBS. Finally, the resulting suspension was adjusted so that the chondrocytes were  
30 suspended in the concentration of 3 x 10<sup>5</sup> cells/ml in the chondrocyte culture medium. The chondrocytes were seeded in an amount of 1 ml/well on 24-well plates. The chondrocytes became confluent 4 days later. The experiment were performed within two weeks after reaching the confluent stage.

##### (b) Addition of Compounds to be Tested and Proteoglycan Depleting Agents

35

The chondrocyte culture medium which had been used for cultivating the chondrocytes was removed from each well and 800 μl of fresh serum-free S-Clone medium containing 0.1% human serum albumin was added. Further, 100 μl of S-Clone medium containing the compounds to be tested (containing the compound in the concentration of 10 fold the final concentration; DMSO concentration = 2.5%) was added.  
40 The chondrocytes were cultured in the presence of carbon dioxide (5%) and air (95%) for 2 hours. Then, the proteoglycan depleting agent, PMA (phorbol myristate acetate) (final concentration = 0.1 μg/ml) or interleukin-1α: (IL-1α) (final concentration = 20 u/ml) was added into the culture medium of the chondrocytes.

The compounds to be tested were as follows:

45

Compounds of present invention: caffeic acid (CFA: Tokyo Kasei Kogyo Co., Ltd.), ferulic acid (FLA: Sigma Chemical Co.), and 3-ethoxy-4-hydroxycinnamic acid (3-EtO-4-HO-CA: compound prepared by Example 1(1))

Comparative substance: Indometacin (Sigma Chemical Co.)

50

##### (c) Determination of proteoglycan

Proteoglycan depletion was determined by the measurement of the glycosaminoglycan (major constituent of proteoglycan, hereinafter referred to as GAG) content following digestion of the chondrocyte matrix with papain.

55

After 2 days, the supernatant of the chondrocyte culture was removed. Then, 1 ml of 0.03% papaine solution was added to the remaining chondrocyte matrix layer and a reaction was performed at 65°C for 1 hour to liberate the GAG from the matrix layer. The content of the GAG in the treated papaine solution was determined by the 1,9-dimethylmethylene blue method (refer to R.W. Farndale, Biochim. Biophys. Acta.,

Vol. 883, pp. 173 to 177, 1986). The GAG content in the chondrocyte matrix of the control test wherein the proteoglycan depleting agent was not added was shown as "100", and the relative amount of the GAG of each experiment except the control test was calculated by the following formula:

$$5 \quad \text{GAG relative amount (\%)} = (B/A) \times 100$$

wherein A represents the GAG content of the control tests wherein the proteoglycan depleting agent was not added, and B represents the GAG content wherein the proteoglycan depleting agents were added alone or the GAG content wherein the proteoglycan depleting agents and the compounds to be tested were added.

The GAG contents of the control tests varied in a range of 28.0 to 74.5  $\mu\text{g/ml}$ , depending on the period from the time when the chondrocytes became confluent until the time when the chondrocytes were used in the above experiment.

The results are shown in Table 2. The GAG content is the value of the mean value  $\pm$  standard error ( $n = 3$ ). For each of the compounds to be tested, the control test and the proteoglycan depleting test wherein the proteoglycan depleting agent was added were carried out and the results thereof are also shown. The significance was determined by Student's t-test with respect to the proteoglycan depleting test wherein the proteoglycan depleting agent was added. The results of the determination are shown as follows:

\*:  $P < 0.05$ ;  
 \*\*:  $P < 0.01$ ;  
 \*\*\*:  $P < 0.001$ .

In comparison with the GAG content in the control tests wherein the proteoglycan depleting agent was not added, the addition of the proteoglycan depleting agents, PMA or IL-1 $\alpha$ , induced a loss of GAG content. Under these conditions, the present compound significantly inhibited or reduced the loss of GAG content, and showed a function to inhibit or suppress the proteoglycan depletion. On the contrary, indomethacin, a conventional analgesic and anti-inflammatory agent, did not show the function to inhibit or suppress the proteoglycan depletion, but showed a significant exacerbation on the cartilage destruction.

Table 2

Samples	GAG content ( $\mu\text{g/ml}$ )	(Relative amount of GAG) (%)
Control	43.2 $\pm$ 1.5**	(100)
IL-1 $\alpha$	33.4 $\pm$ 1.0	(77.3)
IL-1 $\alpha$ + FLA (100 $\mu\text{M}$ )	39.9 $\pm$ 0.8**	(92.4)
IL-1 $\alpha$ + CFA (100 $\mu\text{M}$ )	40.9 $\pm$ 0.7***	(94.7)
Control	43.4 $\pm$ 1.7**	(100)
PMA	32.1 $\pm$ 1.6	(74.0)
PMA + FLA (100 $\mu\text{M}$ )	41.1 $\pm$ 4.2	(94.7)
PMA + CFA (100 $\mu\text{M}$ )	47.7 $\pm$ 1.5***	(110)
Control	74.5 $\pm$ 0.6**	(100)
PMA	46.2 $\pm$ 0.8	(62.0)
PMA + 3-EtO-4-HO-CA (100 $\mu\text{M}$ )	61.5 $\pm$ 0.4***	(82.6)
Control	28.0 $\pm$ 0.7***	(100)
PMA	15.4 $\pm$ 0.5	(55.0)
PMA + indometacin (10 $\mu\text{M}$ )	13.2 $\pm$ 0.6*	(47.1)
(33 $\mu\text{M}$ )	11.7 $\pm$ 0.8**	(41.8)

#### Example 3: Effect of Test Compounds on Proteoglycan Depletion in Mouse Air Pouch Model

The procedure used in the present Example was based on the method described in K.M.K. Bottomley et al., Br. J. Pharmacol., Vol. 93, pp. 627-635 (1988).

The femoral head cartilage (FHC; 40 to 50 mg) was sterily taken out from S.D. rats (5.5 weeks old, female), wrapped in a cotton fabric (5 mm x 5 mm; about 3 mg), and then, implanted subcutaneously in the

backs (where 5 ml of air had been injected the day before) of BALB/c mice (7 weeks old, female). n-Propyl ferulate [nPrFLA; prepared by Example 1(4)] dispersed in an aqueous solution of a mixture of CMC and Tween 80 (final concentration: CMC = 0.3%, Tween 80 = 0.05%) was orally administered at the dose of 100 mg/kg, starting from the third day after implantation, at a rate of five times a week. An aqueous solution of a mixture of the CMC and Tween 80 was orally administered to the control group.

The implanted FHC's were taken out on the 21st day after implantation, and the FHC matrices were digested with a papain solution (4 u/ml papain, 0.05M PBS, pH 6.5, 10 ml). After digestion, the GAG content in the papain solution was determined by the 1,9-dimethyl methylene blue method (R. W. Farndale, vide supra).

In the case of the non-treated group, the FHC's were taken out on the third day after implantation and the GAG content was measured.

The results are shown in Table 3. The GAG content/FHC in the Table 3 is the value of the mean value  $\pm$  standard error (n = 8 to 10) of the GAG content per FHC. The GAG relative amount (%) is the relative amount when the GAG content/FHC of the non-treated group is 100.

Cartilage destructive factor was secreted from the newly formed granuloma around the implanted FHC's. Therefore, a reduction of GAG was observed in the control group. On the contrary, suppression of the GAG reduction was observed in the group to which the present substance was administered and therefore the action of the present substance in suppressing cartilage destruction was confirmed.

Further, methyl ferulate, ethyl ferulate, and n-butyl ferulate exhibited effects similar to that of n-propyl ferulate.

Table 3

Treatment	GAG content/FHC $\mu\text{g}/\text{FHC}$	Relative amount of GAG (%)
No treatment	745.8 $\pm$ 24.5	(100)
Control	387.1 $\pm$ 17.1	(51.9)
nPrFLA	435.3 $\pm$ 26.4	(58.4)

#### Example 4: Formulation of Granule

The following ingredients were mixed homogeneously:

Ferulic acid	20 parts by weight
Lactose	68 parts by weight
Low-substituted hydroxypropylcellulose	10 parts by weight
Hydroxypropylcellulose	2 parts by weight


The mixture was kneaded using 32 parts by weight of a wetting agent, ethanol. Then, the kneaded mixture was granulated by wet granulation and dried to obtain the granule.

As explained above, the present substance strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage. Further, the present substance has low toxicity. Accordingly, the present substance is very useful for the treatment of arthropathy, such as rheumatoid arthritis, osteoarthritis, peri arthritis humeroscapularis, shoulder-arm-neck syndrome, lumbago, and so on.

Although the present invention has been described with reference to specific embodiments, various changes and modifications obvious to those skilled in the art are deemed to be within the spirit, scope, and concept of the invention.

#### Claims

1. A chondroprotective agent comprising a carboxylic acid compound of the general formula (I):



10 wherein R<sup>1</sup> and R<sup>2</sup> are, independently, a hydrogen atom or an alkyl group of 1 to 6 carbon atoms, or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier.

- 15
2. A chondroprotective agent according to claim 1, wherein said ester is an ester with an aliphatic alcohol having 1 to 6 carbon atoms.
  3. A chondroprotective agent according to claim 1, which contains one or more compounds selected from the group consisting of caffeic acid, ferulic acid, isoferulic acid, 3-ethoxy-4-hydroxycinnamic acid, sodium ferulate, methyl ferulate, ethyl ferulate, n-propyl ferulate, and n-butyl ferulate.

20

  4. Use of a compound of the general formula (I), or a cis- or trans-isomer thereof, or a pharmaceutically acceptable salt or ester thereof according to claim 1, for preparing a chondroprotective agent.
- 25
- 30
- 35
- 40
- 45
- 50
- 55



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 94 10 9873

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,X	INDIAN J. EXP. BIOL., vol.25, no.3, March 1987 pages 187 - 189 A.S.CHAWLA ET AL. 'Anti-Inflammatory Action of Ferulic Acid and Its Esters in Carrageenan Induced Rat Paw Oedema Model' * the whole document *	1-4	A61K31/19 A61K31/215
X	ACTA ZOOL. SIN., vol.38, no.1, March 1992 pages 106 - 107 B.-S. ZHOU ET AL. 'ULTRASTRUCTURAL OBSERVATION ON THE EFFECTS OF FERULIC ACID ON THE CARTILAGE PROTEOGLYCANS IN YOUNG RATS' * the whole document *	1-4	
X	WUHAN DAXUE XUEBAO, ZIRAN KEXUEBAN, no.4, 1990 pages 93 - 97 B.ZHOU ET AL. 'ACTION OF DERULIC ACID AND SODIUM SELEMITE ON YOUNG RATS' * abstract *	1-4	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K
X	BIOTECHNOL.THER., vol.3, no.3-4, 1992 pages 119 - 135 B.A. 'T HART ET AL. 'METABOLIC ACTIVATION OF PHENOLS BY STIMULATED NEUTROPHILS: A CONCEPT FOR A SELECTIVE TYPE OF ANTI-INFLAMMATORY DRUG' * the whole document *	1-4	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24 August 1994	Examiner Theuns, H
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 94 10 9873

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	ACUPUNCT. ELECTRO-THER. RES. INT. J., vol.7, no.2-3, 1982 pages 173 - 202 J.CYONG ET AL. 'A PHARMACOLOGICAL STUDY OF THE ANTI-INFLAMMATORY ACTIVITY OF CHINESE HERBS. A REVIEW' * page 179 * * page 185 *	1-4	
A	CLIN.CHIM.ACTA, vol.73, no.1, 1976 pages 121 - 125 M.R.WILLS ET AL. 'THE EFFECT OF URAEMIC METABOLITES ON PARATHYROID EXTRACT-INDUCED BONE RESORPTION IN VITRO' * table I *	1-4	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24 August 1994	Examiner Theuns, H
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document			

EPO FORM 1501 (04/92) (Pec/CO)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**